

LA-UR-96-2643

Title:

A Search for the  $\Delta^+$  Wave-Function Component  
in Light Nuclei

CONF-9605180--5

Author(s):

C. L. Morris, J. D. Zumbro, R. L. Boudrie,  
P. A. M. Gram, J. L. Matthews, Y. Tan,  
V. V. Zelevinsky, E. A. Pusyuk, B. J. Kriss,  
and G. Glass


RECEIVED  
SEP 09 1996  
OSTI

Submitted to:

PANIC'96, XIV International Conference on  
Particles and Nuclei  
May 22-28, 1996  
Williamsburg, Virginia

MASTER

DISTRIBUTION OF THIS DOCUMENT



**Los Alamos**  
NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-10. By acceptance of this article, the publisher recognizes that the U.S. Government retains a non-exclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Form LA 820-06  
81 Rev 10/91



#### **DISCLAIMER**

**Portions of this document may be illegible  
in electronic image products. Images are  
produced from the best available original  
document.**

## A SEARCH FOR THE $\Delta^+$ WAVE-FUNCTION COMPONENT IN LIGHT NUCLEI

C. L. MORRIS, J. D. ZUMBRO, R. L. BOUDRIE, and P. A. M. CHAM,  
*Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

J. L. MATTHEWS, Y. TAN and V. V. ZELEVINSKY  
*Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

E. A. PASYUK  
*Joint Institute for Nuclear Research, Dubna, Moscow Region, 141980 Russia*

B. I. KRISS  
*University of Colorado, Boulder, CO 80309*

G. GLASS  
*9918C Walnut St., Los Alamos, NM 87544, USA*

We have studied the  $(\pi^+, \pi^+ p)$  reactions on  ${}^3\text{H}$ ,  ${}^4\text{He}$ ,  ${}^6\text{Li}$ , and  ${}^7\text{Li}$  at incident energy 500 MeV in quasi-free kinematics. A signature attributable to pre-existing  $\Delta$  components of the ground state wave function is observed.

Meson models of nuclear binding predict both excess pions and excited nucleons in nuclear matter.<sup>1</sup> Results from a new method of measuring the  $\Delta$  component of the nuclear wave function using the pion double charge exchange (DCX) reaction  $(\pi^+, \pi^-)$  are reported in this paper.

If the nucleus contains only nucleons, two like-charge nucleons must participate in the DCX process. If the nuclear wave function contains a  $\Delta^+$ , the  $(\pi^+, \pi^-)$  process can occur in a single step. Recent measurements on a range of nuclei have yielded evidence for this effect.<sup>2</sup> In the current work we have extended these studies to  ${}^3\text{H}$ ,  ${}^4\text{He}$ ,  ${}^6\text{Li}$ , and  ${}^7\text{Li}$ .

The experiment was performed using a 500 MeV beam at the Clinton P. Anderson Meson Physics Facility. Pions from  $(\pi^+, \pi^+ p)$ , NCX, and  $(\pi^+, \pi^- p)$ , DCX, reactions were observed at an angle of 50° in coincidence with protons at 50° using two magnetic spectrometers, in the kinematics of free  $\pi p$  scattering. The sum of the pion and proton energies was required to exceed 400 MeV.

An estimate of the probability for pre-existing  $\Delta$ 's can be obtained from the data by integrating the cross sections and forming the ratio  $R = \sigma_{\text{DCX}}/\sigma_{\text{NCX}}$

$$P_\Delta = R \left( \frac{Z N_\Delta}{A N_\Delta} \right) \left( \frac{\sigma(\pi^+ + p \rightarrow \pi^+ + p)}{\sigma(\pi^+ + \Delta^+ \rightarrow \pi^+ + p)} \right) \left( \frac{k_\Delta}{k_F} \right)^2. \quad (1)$$

Table 1: Measured cross section ratios and extracted values of  $P_{\Delta}$ .

| Target          | $\sigma_{\text{DCX}}/\sigma_{\text{NCX}}$<br>( $\times 10^3$ ) | $P_{\Delta^-}$<br>(%) | $P_{\Delta^+}$<br>(%) |
|-----------------|--|-----------------------|-----------------------|
| ${}^3\text{H}$  | 8.9(6)   | 1.72(12)              | 3.44(24)              |
| ${}^4\text{He}$ | 0.90(18)   | 0.26(5)               | 1.04(20)              |
| ${}^6\text{Li}$ | 0.67(13)   | 0.20(4)               | 0.73(15)              |
| ${}^7\text{Li}$ | 5.4(5)   | 1.68(15)              | 4.22(10)              |

where  $P_{\Delta}$  is the sum of the probabilities for all  $\Delta$  charge states,  $N_{\Delta^-}/N_{\Delta}$  is the fraction of  $\Delta^-$ 's,  $Z$  and  $A$  are the proton and nucleon numbers, and  $k_{\Delta}$  and  $k_F$  characterize the momentum spreads of the  $\Delta$ 's and nucleons. We have used measured branching ratio<sup>3</sup> for  $N^*(1520) \rightarrow \pi + N$  and  $N^*(1520) \rightarrow \pi + \Delta(1232)$  to estimate the cross-section ratio, and have taken  $k_F \approx 200$  MeV/c and  $k_{\Delta} \approx 400$  MeV/c, the peak of the momentum distribution predicted for the calculated pion excess.<sup>4</sup> The ratios of  $\Delta$  charge states have been obtained by generalizing arguments given in<sup>4</sup> to give:

$$\frac{N_{\Delta^-}}{N_{\Delta}} = \frac{3}{4} \left( 1 + \frac{Z}{2(N-1)} + \frac{Z(Z-1)}{N(N-1)} \right)^{-1}, \quad (2)$$

where  $N$  is the neutron number of the nucleus. The measured cross section ratios and resulting  $\Delta$  probabilities are given in Table 1.

The  $\Delta^-$  probability in  ${}^3\text{H}$  is found to agree roughly with theoretical expectations.<sup>5</sup> Moreover, the  $\Delta^-$  probabilities in the light self-conjugate nuclei are approximately an order of magnitude smaller than those in the  $T=1/2$  nuclei. This is presumably because only core-excited two-particle configurations or  $\Delta\Delta$  excitations can contribute to the  $\Delta$  probabilities for neutron pairs coupled to  $J=0$  within the  $s$ -shell.

This work has been supported in part by the U. S. Department of Energy.

#### References

1. B. L. Friman, V. R. Pandharipande, and R. B. Wiringa, *Phys. Rev. Lett.* **51**, 703 (1983).
2. C. L. Morris, J. M. O'Donnell, and J. D. Zumber, *Acta Phys. Pol. B* **24**, 1659 (1993).
3. M. Aguilar-Benitez *et al.*, *Phys. Rev.* **D43S**, 1 (1992).
4. H. J. Lipkin and T.-S. H. Lee, *Phys. Lett. B* **183**, 23 (1987).
5. W. Steune *et al.*, *Nucl. Phys. A* **405**, 661 (1987).